



Randomised controlled trial of inhaled corticosteroids (fluticasone propionate) in cystic fibrosis

Ian M Balfour-Lynn, Nigel J Klein and Robert Dinwiddie

Arch. Dis. Child. 1997;77;124-130
doi:10.1136/adc.77.2.124

Updated information and services can be found at:
<http://adc.bmj.com/cgi/content/full/77/2/124>

These include:

References

This article cites 34 articles, 12 of which can be accessed free at:
<http://adc.bmj.com/cgi/content/full/77/2/124#BIBL>

13 online articles that cite this article can be accessed at:
<http://adc.bmj.com/cgi/content/full/77/2/124#otherarticles>

Rapid responses

You can respond to this article at:
<http://adc.bmj.com/cgi/eletter-submit/77/2/124>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article

Notes

To order reprints of this article go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to *Archives of Disease in Childhood* go to:
<http://journals.bmj.com/subscriptions/>

Randomised controlled trial of inhaled corticosteroids (fluticasone propionate) in cystic fibrosis

Ian M Balfour-Lynn, Nigel J Klein, Robert Dinwiddie

Abstract

Background—Controlling lung inflammation may be the key to improving morbidity and mortality in cystic fibrosis.

Objective—To assess the effects of inhaled corticosteroids on lung inflammation in cystic fibrosis.

Design—Double blind placebo controlled randomised sequence crossover trial. Fluticasone propionate (400 µg/day) was given as a dry powder inhaler for six weeks with a four week washout period before crossover.

Outcome measures—Sputum inflammatory markers (interleukin-8, tumour necrosis factor- α (TNF- α) and neutrophil elastase—both free and bound to α_1 -antiprotease), sputum interleukin-10, lung function, and symptomatology.

Subjects—Twenty three children from a regional cystic fibrosis centre were enrolled into the study, with mean age 10.3 years (range 7 to 17 years) and mean baseline forced expiratory volume in one second (FEV₁) of 64% (range 21% to 102%) predicted for sex and height. One patient was excluded for non-compliance to the study protocol.

Results—No significant benefit was shown for the use of fluticasone propionate in any of the outcomes. For sputum interleukin-8 there was an estimated true treatment median difference of 142 pg/ml (95% confidence interval (CI) 8 to 2866 pg/ml) in favour of placebo; while for maximal expiratory flow at 25% (MEF_{25%}) remaining forced vital capacity predicted for sex and height there was a 15 percentage points (pp) (95% CI 4 to 26 pp) mean treatment difference in favour of placebo. Sputum interleukin-10 was undetected in any samples and unaffected by fluticasone propionate. Neither atopic status, baseline FEV₁, nor concomitant DNase therapy had any effect on response to treatment.

Conclusions—Lack of benefit from fluticasone propionate was most likely due to failure of the drug to penetrate the viscid mucus lining the airways. It is suggested a large multicentre trial with higher doses given for a longer time by a different delivery system is required to assess efficacy.

(*Arch Dis Child* 1997;77:124-130)

Lung disease is responsible for much of the morbidity and most of the mortality associated with cystic fibrosis. Severe lung injury is a result of a continuous cycle of infection and inflammation. Bronchoalveolar lavage (BAL) studies have shown that this inflammation may be present from infancy,^{1,2} is persistent, and is not confined to severe disease or infective exacerbations.³ In some cases, lung inflammation may even precede infection,^{1,4} and it has been suggested that it may be an inherent part of cystic fibrosis.⁵ Consequently, attention has turned increasingly to anti-inflammatory treatment.

Several types of agents have been tried although often only in small trials. Orally active non-steroidal anti-inflammatory drugs such as ibuprofen,⁶ piroxicam,⁷ and pentoxifylline⁸ have been used with variable success. Aerosolised secretory leukoprotease inhibitor⁹ and α_1 -antitrypsin have also been used to combat the excessive neutrophil elastase activity found in the cystic fibrosis lung.¹⁰ Corticosteroids are an obvious choice due to their wide ranging anti-inflammatory effects, particularly on neutrophils, the inflammatory cells which dominate the lung in this disease.¹¹ The initial enthusiasm after the first major trial of oral corticosteroids¹² has been dampened, however, by the follow up report of serious adverse effects.¹³ Since then, a large multicentre trial in the USA recommended that oral corticosteroids should not be continued for more than 24 months, which invalidates this treatment as a long term option.¹⁴

Inhaled corticosteroids would seem the next logical choice due to their localised anti-inflammatory actions in the airways and their relatively high safety profile. Adverse systemic effects are minimal if high doses are avoided and particularly if they are given by spacer device.¹⁵ The primary aim of this study was to assess the effects of inhaled corticosteroids on lung inflammation in children with cystic fibrosis. Inflammation was monitored by sputum interleukin-8, tumour necrosis factor- α (TNF- α), and neutrophil elastase, inflammatory markers, which have all been shown to be raised in cystic fibrosis sputum.^{8,16,17} Another aim was to monitor the effect on sputum interleukin-10, the 'anti-inflammatory' cytokine whose production is downregulated in cystic fibrosis.^{18,19} Finally, lung function and symptomatology were monitored in order to provide pilot data for planning a larger study.

Great Ormond Street
Hospital for Children
NHS Trust,
London: Department
of Respiratory
Medicine
I M Balfour-Lynn
R Dinwiddie

Department of
Immunobiology
N J Klein

Correspondence to:
Dr I M Balfour-Lynn,
Department of Respiratory
Medicine, Great Ormond
Street Hospital for Children
NHS Trust,
Great Ormond Street,
London WC1N 3JH.

Accepted 29 April 1997

Keywords: cystic fibrosis; corticosteroids; inflammation; randomised controlled trial

Methods

SUBJECTS

Children with previously diagnosed cystic fibrosis were enrolled from a single centre. Inclusion criteria were age between 6 and 17 years, ability to reliably perform spirometry and use a Diskhaler (Glaxo Wellcome UK), and guaranteed sputum production. Their routine medication was not altered and no one was started on DNase during the study period. Exclusion criteria were the use of systemic or inhaled corticosteroids within the previous six months, clinical diagnosis of asthma, use of sodium cromoglycate or long acting β_2 -agonists within six weeks, and lower respiratory tract infection requiring antibiotics within the previous three weeks. Other exclusion criteria were cystic fibrosis related diabetes, portal hypertension, allergic bronchopulmonary aspergillosis, or isolation of *Burkholderia cepacia* in sputum. Atopic status was determined by positive personal history of atopy (rhinitis or eczema) as detected by the ISAAC questionnaire,²⁰ or serum IgE > 100 ku/l, or positive skin prick tests to house dust mite, grass pollen, cat and dog fur (presence of at least two weals greater than 3 mm at 20 minutes).

STUDY DESIGN

The trial was conducted in a double blind placebo controlled randomised crossover sequence. After a baseline visit (V0), there were two treatment periods of six weeks' duration with a four week washout period; patients were assessed before and after each period (V1–V4). Each received 400 μ g/day fluticasone propionate in two doses via Diskhaler (dry powder inhaler) or matched placebo provided by Glaxo Wellcome UK. All unused disks were returned and counted to assess patient compliance to the study protocol. The study was approved by the hospital ethics committee; parents or patients, or both, gave written informed consent.

Primary outcome measures were change from baseline in the sputum inflammatory markers interleukin-8, TNF- α , and neutrophil elastase (free and bound to α_1 -antiprotease). Covariates were atopic status and disease severity as assessed by baseline forced expiratory volume in one second (FEV₁). Secondary outcomes were lung function and symptomatology as well as patient tolerance of fluticasone propionate. Adverse events were monitored closely, especially infectious respiratory exacerbations and oral candidiasis.

SPUTUM INFLAMMATORY MARKERS

Sputum was collected into a sterile container, placed onto ice, and weighed. A previously described method was adapted for specimen preparation.²¹ An equal volume of 20% *N*-acetylcysteine (ml for gram) was added to the sputum which was left on ice for 20 minutes. This was then vortexed at 4°C before centrifuging at 4°C for 15 minutes at 3000 \times *g*. The supernatant was decanted and stored in aliquots at –80°C.

After thawing, supernatants were centrifuged at 13000 \times *g* at 4°C for 10 minutes. For

cytokine determination, supernatants were mixed with an equal volume of protease inhibitors (sodium fluoride 20 mM, aprotinin 2%, phenylmethylsulphonyl fluoride 2 mM (Sigma), and 5% human serum known to have undetectable cytokine concentrations). This combination was found to completely inhibit free elastase activity in all sputum samples. Samples were analysed by ELISA for interleukin-8 (using a commercially available kit, Biotrak, Amersham), and TNF- α and interleukin-10 using capture and detector antibodies from PharMingen (San Diego, California, USA), streptavidin-alkaline phosphatase (Amersham) and developed with p-nitrophenyl phosphate disodium (Sigma). Free elastase was measured by a modification of a previously described method.⁸ Sputum supernatants were diluted 1:20 with 0.1M Tris, 0.5 mol/l sodium chloride and 0.05 mmol/l EDTA. This was added to an equal volume of substrate solution (120 μ g *N*-methoxysuccinyl-ALA-ALA-PRO-VAL-p-nitroanilide (Sigma) per ml Tris salt buffer). Optical densities were measured at 405 nm on a Dynatech MRX ELISA plate reader and calibrated using purified human neutrophil elastase (Sigma). Elastase: α_1 -antiprotease complexes were measured by ELISA as previously described.²² After dilution, lower limits of detection were 40 pg/ml for interleukin-8, 30 pg/ml for TNF- α , 250 pg/ml for interleukin-10, 0.05 u/ml for free neutrophil elastase and 1 ng/ml for bound neutrophil elastase.

LUNG FUNCTION

Lung function was assessed by standard spirometry using a compact spirometer (Vitalograph, Buckingham). At each visit, forced vital capacity (FVC), FEV₁, and maximal expiratory flow at 25% remaining FVC (MEF_{25%}) were recorded as per cent predicted for sex and height.²³ Resting oxygen saturation was measured using a Biox 3700e pulse oximeter (Ohmeda, USA). On subsequent clinic visits, lung function was recorded within two hours of the baseline reading and bronchodilators were withheld for four hours before each test.

SYMPTOM SCORES

At the end of each week, the patient or parent recorded symptom scores for cough, sputum production, and wheeze for the previous seven days (combined to provide overall mean respiratory score). At each visit, overall scores for general wellbeing and appetite were recorded. These were indicated on a 10 cm linear analogue scale, which was measured to establish symptom severity.

STATISTICAL METHODS

Analysis was performed by the Applied Statistics Research Unit Ltd at the University of Kent at Canterbury using the SAS System (Release 6.09). Change from baseline for each inflammatory marker was compared using non-parametric Wilcoxon rank sum test for crossover design (Koch's method).²⁴ Per cent predicted lung function at the end of each

Table 1 Baseline lung function (mean % predicted with range) and mean coefficient of variation (%) with adjusted mean (95% CI) at end of treatment with fluticasone propionate and placebo. Also included are estimated mean treatment differences (fluticasone propionate minus placebo) with 95% CI. Values of $p > 0.05$ are considered non-significant (NS)

	FVC (%)	FEV ₁ (%)	MEF _{25%} (%)	Oxygen saturation (%)
Baseline	75 (45 to 107)	64 (21 to 102)	35 (8 to 73)	95 (90 to 98)
Coefficient of variation (%)	7	9	24	–
Fluticasone propionate	78 (74 to 83)	65 (61 to 69)	25 (15 to 36)	95.0 (94.6 to 95.3)
Placebo	78 (73 to 82)	67 (63 to 71)	40 (30 to 50)	95.4 (95.0 to 95.8)
Estimated mean treatment difference	0.5 (–6 to 7)	–2 (–8 to 4)	–15 (–26 to –4)	–0.4 (–1.0 to 0.1)
p Value	NS	NS	0.009	NS

Table 2 Baseline sputum inflammatory markers (median with interquartile range) and mean coefficient of variation (%) with median (95% CI) change from baseline after treatment with fluticasone propionate and placebo. Also included are estimated median treatment differences (fluticasone propionate minus placebo) with 95% CI. Values of $p > 0.05$ are considered non-significant (NS)

	Interleukin-8 (pg/ml)	TNF- α (pg/ml)	Free neutrophil elastase (u/ml)
Baseline	376 (306 to 437)	375 (198 to 583)	62 (40 to 80)
Coefficient of variation (%)	54	30	26
Fluticasone propionate	8 (–161 to 217)	14 (–90 to 103)	–1 (–8 to 14)
Placebo	–67 (–1152 to 37)	25 (–122 to 117)	1 (–15 to 7)
Estimated median treatment difference	142 (8 to 2866)	27 (–104 to 167)	5 (–9 to 31)
p Value	0.03	NS	NS

treatment, and change from baseline in symptom scores were compared using parametric analysis of covariance. The covariates used were atopic status, baseline per cent predicted FEV₁, and colonisation with *Pseudomonas aeruginosa*. In addition, for FVC and MEF_{25%} variables, their mean baseline values were also used. Values from the run-in and washout period were used to detect any first order and second order carryover effects. If carryover was detected, analysis was based on the first treatment period only. Significance was judged at the 10% level for carryover effects and at the 5% level for treatment and period effects. All statistical tests were two sided. Effect of atopic status and use of DNase on treatment difference in change from baseline of inflammatory markers was assessed using subgroup analyses and employing the non-parametric method of Koch; while effect on treatment difference in end of treatment lung function was assessed by analysis of covariance. Pre-existing lung function was calculated using baseline per cent predicted FEV₁ (mean of FEV₁ measured at V0 and V1). For the inflammatory markers, regression analysis was performed plotting baseline FEV₁ against the log ratio before and after treatment levels for each treatment, and for lung function, baseline FEV₁ was plotted against after treatment lung function. Retrospective sample size calculations for crossover studies based on the observed variability were performed for a power of 80% and a 5% significance level.

Results

Twenty three patients were enrolled into the study and randomised, but one patient failed to comply with treatment so the efficacy sample used for analysis consisted of 22 patients. The two sequence groups were comparable in terms of demographic and baseline clinical characteristics. There were 11 boys and 12 girls with a mean age of 10.3 years (range 7–17 years). Fifteen (65%) were homozygous for $\Delta F508$ gene mutation, seven (30%) were heterozygous for $\Delta F508$, and one child was heterozygous for G551D mutation. Mean (SD) centiles pre-

dicted for sex and age were 40% (36%) for weight and 43% (34%) for height. Table 1 shows mean baseline per cent predicted lung function. Seventeen (74%) patients reported daily cough and 22 (96%) daily sputum production; seven (30%) patients wheezed, although only two on a daily basis. Three were bronchodilator responsive (increase in FEV₁ >15% after inhaled β -agonist), none of whom reported wheezing. Ten (43%) patients were considered atopic. Previous sputum microbiology revealed 96% had been infected with *Staphylococcus aureus*, 87% with *P aeruginosa*, and 48% with *Haemophilus influenzae*. At the start of the study, 61% had positive sputum culture for *S aureus*, 52% for *P aeruginosa*, and 4% (one patient) for *H influenzae*. Oral antibiotics were started in 18 (78%) patients during the trial and nebulised antibiotics started in five (22%). Five (22%) patients required intravenous antibiotics. Antibiotic usage was the same for patients in both treatment sequences. Nine (39%) patients were on nebulised DNase at the start of the trial and continued its use; none started DNase during the trial.

SPUTUM INFLAMMATORY MARKERS

Table 2 shows sputum interleukin-8, TNF- α , and free neutrophil elastase baseline levels (taken at V1) with median change after treatment with fluticasone propionate and placebo. Figure 1 shows treatment changes for individual patients. For interleukin-8, estimated true treatment median difference (fluticasone propionate–placebo) was 142 pg/ml in favour of placebo (with 95% confidence interval (CI) for difference of 8 to 2866 pg/ml). This was a statistically significant difference with $p=0.03$. There was little change from baseline for sputum TNF- α or free neutrophil elastase during either fluticasone propionate or placebo, with no significant difference between the two treatments. In 16 out of 23 patients, at baseline, bound neutrophil elastase was undetectable (<10 ng/ml), in the others, it ranged from 2–42 (median 10) ng/ml. Analysis of change in bound neutrophil elastase is invalid




Figure 1 Sputum inflammatory markers: (A) interleukin-8 (pg/ml), (B) TNF- α (pg/ml), and (C) free neutrophil elastase (u/ml) before (solid circles) and after (open circles) treatment with fluticasone propionate and placebo. Median values with 95% CI are also shown (bars and vertical lines). Y axes for interleukin-8 and TNF- α are on log scale.

due to the large number of samples with undetectable levels (80/110). Neither atopic status, baseline FEV₁, nor concomitant DNase therapy had any significant effect on response to treatment. An estimate of coefficient of variation for the markers was calculated using values from the initial visit (V0) and the first visit (V1) before treatment. Table 2 shows that these were highest for interleukin-8. Sputum interleukin-10 was undetectable in all samples; no effects were seen after giving corticosteroids.

LUNG FUNCTION

No effect of treatment was seen on either FVC or FEV₁ (see table 1). For MEF_{25%} there was a significant second order carryover effect ($p=0.02$) so analysis was based on the first treatment period only (12 patients on flutica-

sone propionate, 10 on placebo). Adjusted mean per cent predicted MEF_{25%} was higher after treatment with placebo than fluticasone propionate (table 1). The estimated difference between the treatments (fluticasone propionate - placebo) was -15pp (95% CI -26 to -4 pp) which was statistically significant with $p=0.009$. Resting oxygen saturation was unaffected by either treatment (table 1). Neither atopic status, baseline FEV₁, nor concomitant DNase therapy had any significant effect on response to treatment. From the measurements at V0 and V1, mean coefficient of variation for lung function was highest for MEF_{25%} (see table 1).

SYMPTOM SCORES

Table 3 shows baseline symptom scores with mean changes after treatment. Mean overall respiratory score, wellbeing, and appetite scores were all unaffected by treatment.

ADVERSE EVENTS

Two patients (9%) experienced an adverse event which was possibly drug related during treatment with fluticasone propionate. One child coughed immediately after taking fluticasone propionate for five minutes on one occasion, while another developed peeling skin on the fingers for three days, the cause of which was unknown. Overall 18 (78%) patients had adverse events during treatment with fluticasone propionate and 15 (65%) while on placebo. These were mainly upper respiratory tract infections (52%), or cystic fibrosis chest exacerbations (20%).

POWER STUDIES

Retrospective sample size estimation for crossover studies have been calculated, based on the variability observed in this study. For change from baseline in sputum inflammatory markers the sample size estimation assumed the data were normally distributed, which was not actually the case, even after taking a log_e transformation. For interleukin-8, to detect a difference of 150 pg/ml, it was estimated that huge numbers of patients were needed. In fact, the huge variability of interleukin-8 was due to a small number of patients and furthermore, with non-parametric analysis a statistically significant difference in interleukin-8 was obtained in our study. To obtain a difference of 125 pg/ml TNF- α , 46 patients would be needed, and for a difference of 20 u/ml free neutrophil elastase, 18 patients. To obtain a difference of 15% improvement over placebo in per cent predicted FVC, FEV₁, and MEF_{25%}, a study would need 28, 42, and 98 patients respectively. To detect a treatment difference in mean symptom scores of 1.5 requires 14 patients.

Discussion

This study has not shown any significant benefit in terms of sputum inflammatory markers, lung function, or symptomatology when an inhaled corticosteroid (400 μ g/day fluticasone propionate) was given by dry powder inhaler to children for six weeks. However,

Table 3 Baseline symptom scores out of a maximum of 10 points (mean with range) with adjusted mean (95% CI) change from baseline at end of treatment with fluticasone propionate and placebo. Also included are estimated mean treatment differences (fluticasone propionate minus placebo) with 95% CI. Values of $p > 0.05$ are considered non-significant (NS)

	Respiratory	Wellbeing	Appetite
Baseline	3.4 (0.9 to 5.9)	6.7 (3.2 to 9.3)	6.7 (0.1 to 9.4)
Fluticasone propionate	0.5 (-0.3 to 1.2)	-0.4 (-1.0 to 0.2)	0.1 (-1.3 to 1.5)
Placebo	-0.3 (-1.0 to 0.4)	-0.6 (-1.1 to 0)	-0.8 (-2.2 to 0.6)
Estimated mean treatment difference	0.8 (-0.3 to 1.8)	0.1 (-0.7 to 1.0)	0.9 (-0.6 to 2.4)
p Value	NS	NS	NS

with all the theoretical reasons why inhaled corticosteroids should be beneficial in cystic fibrosis,¹¹ the disappointing result needs to be explained, and there are a number of possibilities. Firstly, there may have been a problem with the outcome measures used. The primary aim was to determine the effect on lung inflammation, so inflammatory markers were measured in the sputum which is non-invasive and easy to collect. Sputum markers can give highly variable results, however, since sputum is a non-homogeneous material²⁵ and each cough produces a sample from a different part of the lung. Cystic fibrosis sputum is also a difficult substance to work with in the laboratory due to its viscosity. Furthermore, a recent longitudinal study has shown that sputum cytokine concentrations can vary several fold in the same patient over time, without discernible change in clinical status or lung function.²⁶ Few studies have found correlations between cytokine concentrations and clinical status²⁷ and data on the effect of antibiotics on sputum cytokines is limited.²⁸ Although this throws into doubt the clinical use of sputum markers, it does not invalidate their use in a trial of anti-inflammatory treatment where inflammation is the primary outcome.

With the difficulties in using sputum, many studies, particularly in the USA, measure markers in BAL fluid. BAL is invasive, however, and this would be a problem in a study requiring repeated samples. BAL also has methodological problems, particularly related to the effects of dilution and variability.²⁹ Numerous studies have looked for surrogate markers of inflammation in blood. Aside from the problem of repeated venepuncture in children, none have been shown to be highly predictive of acute changes, and there is often a wide overlap between levels found in health and disease.²⁹ Besides, cytokines tend to act locally in low concentrations so their direct measurement in tissue fluids from the site of production is preferable to measurement of blood levels, which often do not reflect concomitant changes found in sputum or BAL fluid.^{30 31}

Aside from the concerns over sputum, it is also possible there was a problem with the actual marker substances even though they had all been previously shown to be raised in cystic fibrosis.^{8 16 17 21 26 30-32} The levels of TNF- α from this study are in general agreement with those found by others.^{21 26 31} The interleukin-8 levels on the other hand varied widely, both between and within studies with large ranges obtained.^{16 26 30} In this study it can be seen that the majority of samples fell within a fairly nar-

row band, although in some patients huge differences were found for interleukin-8. This may explain the unexpected and possibly random finding that placebo was superior to fluticasone propionate in reducing sputum interleukin-8 levels. There was also a large variability between the two baseline samples, and the reason this was so high compared with that found with the TNF- α , may be due to the less restricted range of cells capable of producing interleukin-8.³³ This may artefactually increase interleukin-8 concentrations in some sputum samples, for example, in the presence of a large number of epithelial cells. Results from this and other studies indicate caution is needed before using sputum interleukin-8 as an inflammatory marker in cystic fibrosis.

The fact that FVC and FEV₁ were unaffected by six weeks of corticosteroids is not surprising, given the fluctuating nature of cystic fibrosis lung disease. Although four weeks is long enough to detect an effect of fluticasone propionate in asthmatic children,³⁴ the nature of cystic fibrosis means it is likely a longer study would be necessary to detect an effect on lung function. The baseline variability for FVC and FEV₁ was reasonable in our study, particularly when compared with the known variability of cystic fibrosis lung function.^{35 36} The variability of MEF_{25%} was high, but it is known to be an intrinsically more unstable measurement. This, together with the fact that due to carryover effect on MEF_{25%} only the first treatment sequence was analysed, may explain why the MEF_{25%} was unexpectedly (and possibly randomly), significantly higher after placebo compared with fluticasone propionate.

It is possible there were too few patients in this trial, although it was as large as most studies of this nature. The problem arose because of the variability of the outcomes used, as discussed above. The variability was apparent once the trial was completed, and it means large numbers of patients are needed unless large treatment differences are found, as shown by our power calculations. Many potential subjects were ruled out by the need to guarantee a sample of sputum each visit, even when well. Recruitment difficulties were also due to the number of visits required for the trial as many children did not want to miss school. We also found that almost 20% of our patient population were already taking inhaled corticosteroids to control troublesome wheezing. This study has highlighted the difficulties in carrying out intervention studies in a disease where the patients are already overburdened by their treatment regimens, and emphasises the need for multicentre trials in cystic fibrosis.

It was also possible that it was already too late for the children in this trial to benefit from inhaled corticosteroids. Lung inflammation is seen early in cystic fibrosis^{1 2} and damage to the lung tissue is relentless. Although most of the subjects studied here had reasonable lung function (mean FEV₁ 64% predicted), there was a wide range (FEV₁ 21% to 102% predicted), and indeed three patients had FEV₁ <40% predicted. In addition, they had to be regular sputum producers to enter the trial and this would bias the subjects towards those with worse lung disease. Not surprisingly, trials of inhaled corticosteroids restricted to adults have so far also been negative.^{37 38} It is likely that maximum benefit of anti-inflammatory therapy will only be seen when treatment is started early, maybe even at the time of diagnosis. The problem will be proving this benefit, as long term studies will be needed and outcome measures for infants with cystic fibrosis are notoriously difficult to use.²⁹

The likeliest reason, however, for the failure to show benefit from inhaled corticosteroids was that the drug did not penetrate the thick viscid sputum. This would reduce the steroid effect on the inflammatory cells contained in the sputum as well as the underlying respiratory epithelial cells. This is backed up by the fact that systemic corticosteroids seem to exert an anti-inflammatory effect in cystic fibrosis^{12 14 39} whereas inhaled corticosteroids have so far not proved to be of benefit. The first trial of inhaled corticosteroids, in 26 subjects, used a low dose (400 µg/day beclomethasone dipropionate for 16 weeks), and showed no beneficial effect on lung function or inflammatory markers.⁴⁰ The next study, over 10 years later, on 12 adults (1600 µg/day budesonide for six weeks), showed an improvement in bronchial responsiveness only, with no change in lung function.³⁷ Another study, on 49 hospitalised patients (1500 µg/day beclomethasone dipropionate for 30 days), showed an improvement over those on standard treatment alone in terms of thoracic gas volume only.⁴¹ Finally, a recently reported abstract on 36 adults (1000 µg/day fluticasone propionate for up to 24 months) showed a non-significant trend only towards improvement in lung function.³⁸

A dry powder inhaler was used in this study to facilitate checking of patient compliance; normally inhaled corticosteroids would be given by a metered dose inhaler with a spacer to reduce oral absorption. It is possible that corticosteroids would penetrate the tenacious sputum more effectively if the drug was delivered as a droplet by spacer or nebuliser, but this is only speculation. A further trial of inhaled corticosteroids is still warranted using a different delivery system. Unfortunately, there would most likely be problems with compliance if a new twice daily nebulised treatment was added to a patient's existing regimen, particularly if no immediate benefit was felt. It is also possible that a higher dose should be used to counter the problems of penetrating the mucus. The dose of 400 µg/day fluticasone propionate used in this study was already above the current licensed dose. It is likely that at higher doses

adrenal suppression would be seen in some patients but in a disease such as cystic fibrosis it may be acceptable to take increased risks as long as these are accompanied by major benefits. Besides, side effects should still be minimal compared with those experienced after long term oral corticosteroids.

In conclusion, controlling lung inflammation may be the key to improving morbidity and mortality in cystic fibrosis, but long term anti-inflammatory treatment needs to be safe and acceptable to the patients. Using inhaled corticosteroids in children with cystic fibrosis is still a good idea in theory. Unfortunately, in common with others, we were unable to prove this during the present trial. Evidence is still needed before too many patients are given this form of treatment on empirical grounds, thus making further trials difficult to conduct. The main problem would seem to be in getting adequate doses of the drug into the relevant cells. The next stage should be a large multicentre trial with higher doses given for a longer time using a different delivery system.

We would especially like to thank the children and their parents for participating in this study. We would also like to thank Dr S Carr and Ms S Madge for their help with specimen collections and Ms T Maslen for help with statistical analysis.

IBL received financial support from Glaxo Wellcome UK.

- 1 Khan TZ, Wagener JS, Bost T, Martinez J, Accurso FJ, Riches DWH. Early pulmonary inflammation in infants with cystic fibrosis. *Am J Respir Crit Care Med* 1995;151:1075-82.
- 2 Armstrong DS, Grimwood K, Carzino R, Carlin JB, Olinsky A, Phelan PD. Lower respiratory tract infection and inflammation in infants with newly diagnosed cystic fibrosis. *BMJ* 1995;310:1571-2.
- 3 Konstan MW, Hilliard KA, Norvell TM, Berger M. Bronchoalveolar lavage findings in cystic fibrosis patients with stable, clinically mild lung disease suggest ongoing infection and inflammation. *Am J Respir Crit Care Med* 1994;150:448-54.
- 4 Balough K, McCubbins M, Weinberger M, Smits W, Ahrens R, Fick R. The relationship between infection and inflammation in the early stages of lung disease from cystic fibrosis. *Pediatr Pulmonol* 1995;20:63-70.
- 5 Cantin A. Cystic fibrosis lung inflammation: early, sustained and severe. *Am J Respir Crit Care Med* 1995;151:939-41.
- 6 Konstan MW, Byard PJ, Hoppel CL, Davis PB. Effect of high-dose ibuprofen in patients with cystic fibrosis. *N Engl J Med* 1995;332:848-54.
- 7 Sordelli DO, Macri CN, Maillie AJ, Cerquetti MC. A preliminary study on the effect of anti-inflammatory treatment in cystic fibrosis patients with *Pseudomonas aeruginosa* lung infection. *Int J Immunopathol Pharmacol* 1994;7:109-17.
- 8 Aronoff SC, Quinn FJ, Carpenter LS, Novick WJ. Effects of pentoxifylline on sputum neutrophil elastase and pulmonary function in patients with cystic fibrosis: preliminary observations. *J Pediatr* 1994;125:992-7.
- 9 McElvaney NG, Nakamura H, Birrer P, et al. Modulation of airway inflammation in cystic fibrosis. In vivo suppression of interleukin-8 levels on the respiratory epithelial surface by aerosolization of recombinant secretory leukoprotease inhibitor. *J Clin Invest* 1992;90:1296-301.
- 10 McElvaney NG, Hubbard RC, Birrer P, et al. Aerosol α1-antitrypsin treatment for cystic fibrosis. *Lancet* 1991;337:392-4.
- 11 Balfour-Lynn IM, Dinwiddie R. Role of corticosteroids in cystic fibrosis lung disease. *J R Soc Med* 1996;89(suppl 27):8-13.
- 12 Auerbach HS, Williams M, Kirkpatrick JA, Cotten HR. Alternate day prednisolone reduces the morbidity and improves pulmonary function in cystic fibrosis. *Lancet* 1985;ii:686-8.
- 13 Donati MA, Haver K, Gerson W, Klein M, McLaughlin FJ, Wohl MEB. Long term alternate day prednisolone therapy in cystic fibrosis. *Pediatr Pulmonol Suppl* 1990;5:277.
- 14 Eigen H, Rosenstein BJ, FitzSimmons S, Schidlow DV, Cystic Fibrosis Foundation Prednisone Trial Group. A multicenter study of alternate-day prednisone therapy in patients with cystic fibrosis. *J Pediatr* 1995;126:515-23.
- 15 Russell G. Inhaled corticosteroid therapy in children: an assessment of the potential for side effects. *Thorax* 1994;49:1185-8.
- 16 Richman-Eisenstat JBY, Jorens PG, Hébert CA, Ueki I, Nadel JA. Interleukin-8: an important chemoattractant in sputum of patients with chronic inflammatory airway

- diseases. *Am J Physiol* 1993;**264**(Lung Cell Mol Physiol 8):L413-8.
- 17 O'Connor CM, Gaffney K, Keane J, et al. α_1 -Proteinase inhibitor, elastase activity, and lung disease severity in cystic fibrosis. *Am Rev Respir Dis* 1993;**148**:1665-70.
 - 18 Bonfield TL, Panuska JR, Konstan MW, et al. Inflammatory cytokines in cystic fibrosis lungs. *Am J Respir Crit Care Med* 1995;**152**:2111-8.
 - 19 Bonfield TL, Konstan MW, Burfiend P, Panuska JR, Hilliard JB, Berger M. Normal bronchial epithelial cells constitutively produce the anti-inflammatory cytokine interleukin-10, which is downregulated in cystic fibrosis. *Am J Respir Cell Mol Biol* 1995;**13**:257-61.
 - 20 Asher MI, Keil U, Anderson HR, et al. International study of asthma and allergies in childhood (ISAAC): rationale and methods. *Eur Respir J* 1995;**8**:483-91.
 - 21 Greally P, Hussein MJ, Cook AJ, Sampson AP, Piper PJ, Price JF. Sputum tumour necrosis factor- α and leukotriene concentrations in cystic fibrosis. *Arch Dis Child* 1993;**68**:389-92.
 - 22 Finn A, Morgan P, Rebuck N, et al. Effects of inhibition of complement activation using recombinant soluble CR1 on neutrophil CD11b/CD18 and L-selectin expression and release of IL8 and elastase in simulated cardiopulmonary bypass. *J Thorac Cardiovasc Surg* 1996;**112**:451-9.
 - 23 Rosenthal M, Bain SH, Cramer D, et al. Lung function in white children aged 4 to 19 years: I—spirometry. *Thorax* 1993;**48**:794-802.
 - 24 Koch GG. Use of non-parametric methods in the statistical analysis of the two-period change-over design. *Biometrics* 1972;**28**:577-84.
 - 25 Cantin AM, Escher E. Protease inhibitors. *Pediatr Pulmonol Suppl* 1994;**10**:159-60.
 - 26 Salva PS, Doyle NA, Graham L, Eigen H, Doerschuk CM. TNF- α , soluble ICAM-1, and neutrophils in sputum of cystic fibrosis patients. *Pediatr Pulmonol* 1996;**21**:11-9.
 - 27 Richman-Eisenstat J. Cytokine soup: making sense of inflammation in cystic fibrosis. *Pediatr Pulmonol* 1996;**21**:3-5.
 - 28 Dean TP, Shute JK, Briars G, Connett G, Church MK, Warner JO. Effect of antibiotic therapy in cystic fibrosis (CF) on sputum concentrations of inflammatory markers. *Eur Respir J* 1995;**8** (suppl 19):511S.
 - 29 Ramsey BW, Boat TF. Outcome measures for clinical trials in cystic fibrosis. Summary of a cystic fibrosis foundation consensus conference. *J Pediatr* 1994;**124**:177-92.
 - 30 Dean TP, Dai YD, Shute JK, Church MK, Warner JO. Interleukin-8 concentrations are elevated in bronchoalveolar lavage, sputum, and sera of children with cystic fibrosis. *Pediatr Res* 1993;**34**:159-61.
 - 31 Kronborg G, Hansen MB, Svenson M, Fomsgaard A, Hoiby N, Bendtzen K. Cytokines in sputum and serum from patients with cystic fibrosis and chronic *Pseudomonas aeruginosa* infection as markers of destructive inflammation in the lungs. *Pediatr Pulmonol* 1993;**15**:292-7.
 - 32 Konstan MW, Berger M. Infection and inflammation of the lung in cystic fibrosis. In: Davis PB, ed. *Cystic fibrosis*. New York: Marcel Dekker Inc, 1993.
 - 33 Adams DH, Lloyd AR. Chemokines: leucocyte recruitment and activation cytokines. *Lancet* 1997;**349**:490-5.
 - 34 MacKenzie CA, Weinberg EG, Tabachnik E, Taylor M, Havnen J, Crescenzi K. A placebo controlled trial of fluticasone propionate in asthmatic children. *Eur J Pediatr* 1993;**152**:856-60.
 - 35 Nickerson BG, Lemen RJ, Gerdes CB, Wegmann MJ, Robertson G. Within-subject variability and per cent change for significance for spirometry in normal subjects and in patients with cystic fibrosis. *Am Rev Respir Dis* 1980;**122**:859-68.
 - 36 Cooper PJ, Robertson CF, Hudson IL, Phelan PD. Variability of pulmonary function tests in cystic fibrosis. *Pediatr Pulmonol* 1990;**8**:16-22.
 - 37 Van Haren EHJ, Lammers J-WJ, Festen J, Heijerman HGM, Groot CAR, van Herwaarden CLA. The effects of the inhaled corticosteroid budesonide on lung function and bronchial hyperresponsiveness in adult patients with cystic fibrosis. *Respir Med* 1995;**89**:209-14.
 - 38 Nieman R, Williams S, Maden C, Knight R, Hodson M. A double-blind placebo-controlled study comparing the effects of the inhaled corticosteroid fluticasone propionate 500 mcg BID with placebo in adults with cystic fibrosis. *Am J Respir Crit Care Med* 1996;**153**:A72.
 - 39 Greally P, Hussain MJ, Vergani D, Price JF. Interleukin-1 α , soluble interleukin-2 receptor, and IgG concentrations in cystic fibrosis treated with prednisolone. *Arch Dis Child* 1994;**71**:35-9.
 - 40 Schiøtz PO, Jørgensen M, Flensborg EW, et al. Chronic *Pseudomonas aeruginosa* lung infection in cystic fibrosis. *Acta Paediatr Scand* 1983;**72**:283-7.
 - 41 Nikolaizik WH, Schöni MH. Pilot study to assess the effect of inhaled corticosteroids on lung function in patients with cystic fibrosis. *J Pediatr* 1996;**128**:271-4.